Appl. Ent. & Phytopath.

Vol. 57. Nos: 1 & 2, Feb. 1990

Summary III bet augment somibendation of the Aret W. MAINAWIND

GREEN OAK LEAF - ROLLER MOTH IN KOHKILUYEH AND BOYER - AHMAD PROVINCE

ABAT M. & K. ATVILL 1996, Vertex consider L. in Fran. Occi. Forest and

(Tortrix viridana L., LEP : TORTRICIDAE)

M. J. FAZELI & M. ABAI

Plant Pests & Diseases Research Institute

Tortrix viridana L., an important species of Tortricidae is considered as a serious pest of the forests of the European countries and the whole U. S. S. R., north - western Africa, and the Near East.

In Iran it was first observed in Yasoudj - in 1983 by Abai. Now it covers all the forests up to Kam - firouz and Chahar - Mahal - Bakhtiari provinces in Iran .

A moth with a wing span of about 18mm., the forewing bright green and hind - wing greyish. The larva which attains a fully grown length of about 15 mm. and is greenish grey with black spots and a black head.

It has one gerneration per year. Winter is passed as egg, laid singly or in pairs, on the young braches.

The neonates penetrate in the buds. The larvae become full - grown until the begining of May, and then pupate inside the group of leaves stick together. The first moth appears around the second half of May.

Bactospein and Dimilin have been tested against the pest, while the second one has shown satisfactory results.

References

- ABAI, M, & E. ADELI, 1984. List of Pests of Forest Trees and Shrubs of Iran,
 Plant Pests and Diseases Research Institute, Tehran, pp: 89 100.
- ABAI, M. & E. ADELI, 1986. *Tortrix viridana* L. in Iran, Dept. Forest and rangland of Min. of Agric. Chalus, 6pp.
- SABETI, H., 1976. Forest trees and shrubs of Iran, Agriculture & Natural Resources Research Organization, Ministry of Agriculture, Tehran.
- SCHWENKE, W. 1978. Die Forestschaedlinge Europas, Bd. III. Schmetter linge, Paul Parey, Hamburg. pp: 76 85.
- WILTSHIRE, E. P. 1957. The Lepidoptera of Iraq, Min. Agric. Baghdad, 55 pp.

Address of the authors:

Eng. M. J. FAZELI. Plant Pests & Diseases Research Department.

Agriculture and Natural Resources Research Centre for Shiraz. P. O. Box :781,
Shiraz 71 365, Iran.

Dr. M. ABAI. Research Department for Harmful Insects and Animals to Plants. Plant Pests & Diseases Research Institute. P. O. Box: 1454, Tehran - 19395, Iran

at and is greenish grey with black spots and a black head.

irs, on the young braches.

The neonates penetrate in the buds. The larvae become full - grown until

shirst moth appears around the second half of May.

Bacrospein and Dimilia have been tested against the pest, while the