Appl. Ent. Phytopath and sall of anotheliugon authorized Discommon Mandallian Vol. 64, 1&2, Feb. 1997. Ison noticed on patients and allowed Barrier and Indiana.

MONOSTON Possible presence of satellite RNA in the AMALETTIMAD

Y. FAZLALI, A. AHOONMANESH, M. R. HAJIMORAD AND A. KARIMI
College of Agriculture, Tarbiat Modarres University, Tehran, College of Agriculture,
Isfahan University of Technology, Isfahan, Plant Petes and Diseases Research

KEARNEY, C. M.; ZIETER, T. A. TOARTSBA. VES, D. (1990). A field survey for

Thirty five isolates of Cucumber Mosaic Virus (CMV) from a number of plant species (mainly tomato) collected from different parts of Iran were tested for the presence of satellite ribonucleic acids (Satellite RNA).

The isolates propagated in tobacco were subjected to dot-blot hybridization (Nicotiana tabacum cv. Havana). Extracts from systemically infected and healthy plants were placed on nylon membranes and sent to Italy and U.S.A. All the samples were checked by CMV-S strain having satellite RNA according to Crescenzi et al., as positive control.

Absence of satellite- RNAs in Iranian isolates under confirms the point that the field conditions in Iran are favour of virus but is not ideal for stimulation and propagataion of satellite RNAs of CMV.

The authors of This paper take this opportunity to thank Dr. D. Gallitelli from Italy and Dr. M. T. Roossinck from U.S.A. for having done "Dot-Blot Nucleic Acid Hybidization Assay" Pertaining to this paper.

References

COLLMER, C. W. & HOWELL, S. H. 1992. Role of satellite RNA in the expression of syptoms by plant viruses. Ann. Rev. of Phytopathol. 30,419-442.

- CRESCENZI, A.; BARBAROSSA, L.; GALLITELLI, D. & MARTELLI, G. P. 1993.

 Cucumber Mosaic Cucumovirus populations in Italy under natural epidemic conditions and after a satellite mediated protection test. Plant Disease 71,28-33.
- FRANCKI, R. I. B. 1985. Plant Virus Satellites. Ann. Rev. of Phytopathol. 39,151-174.
- GALLITELLI, D.; VOVLAS, C.; MARTELLI, G.; MONTASSER, M. S.; TOUSIGNANT, M. E. & KAPER, J. M. 1991. Satellite-mediated protection of tomato against Cucumber Mosaic Virus: II. Field test under natural epidemic conditions in southern Italy. Plant Disease 75,93-95.
- HIBI, T. & SAITO, Y. 1985. A dot immunobinding assay for the detection of Tobacco Mosaic Virus in infected tissues. J. Gen. Virology 66,1191-1194.
- KAPER, J. M. & COLLMER, C. W. 1988. Modulation of viral plant diseases by secondary RNA agents. In: Domingo *et al.* RNA genetics, CRC Press.
- KEARNEY, C. M.; ZITTER, T. A. & GONSALVES, D. (1990). A field survey for serogroups and satellite RNAs of cucumber mosaic virus. Phytopathology 80,1238-1243.
- LIU, Y. Y. & COOPER, J. I. 1994. satellites of plant viruses. Review of Plant
 Pathology 73,371-387.
- MORIONES, E.; DIAZ, I.; RODRIGUEZ-CREZO, E.; FRAILE, A. & GARCIA-ARENAL, F. 1992. Differential interactions among strains of Tomato Aspermy Virus and satellite RNA of Cucumber Mosaic Virus. Virology 186,475-480.
- MORIONES, E.; FRAILE, A. & GARCIA-ARENAL., F. 1991. Host-associated selection of sequence variants from a satellite RNA of Cucumber Mosaic Virus.

 Virology 184,465-468.
- MURANT, A. F. & MAYO, M. A. 1982. Satellites of plant viruses. Ann. Rev. of Phytopathol. 20,49-70. ob galved to A. & U mont stonies on M. T. M. 'I'll bas
- PALUKAITIS, P.; COTTS, S. & ZAITLIN, M. 1985. Detection and identification of viroids and viral nucleic acids by dot-blot hybridization. Acta Horticulture 164,109-118.
- PALUKAITIS, P.; ROSSINCK, M. J.; DITZGEN, R. G. & FRANCKI, R. I. B. 1992.

 Cucumber Mosaic Virus. Advances in Virus Research 41,281-348.

- PIAZZOLLA, P.; DIAZ-RUIZ, J. R. & KAPER, J. M. 1979. Nucleic acid homologies of eigtheen Cucumber Mosaic Virus isolates determined by competition hybridization. J. Gen. Virology 45,361-369.
- ROOSSINCK, M. J.; SLEAT, D. E. & PALUKAITIS, P. 1992. Satellite RNAs of plant viruses: Structures and biological effects. Microbiological Review. 56,265-279.
- VAN REGENMORTEL, M. H. V. 1982. Serology and Immunochemistry of plant viruses. Academic press.
- Address of the authors: Eng. Y. FAZLALI¹, Dr A. AHOONMANESH², Dr M. R. HAJIMORAD AND Eng. A. KARIMI³

 College of Agriculture, Tarbiat Modarres University, Tehran¹; College of Agriculture; Isfahan University of Technology, Isfahan²; Plant Pests and Diseases Research Institute, P. O. Box 1454, 19395, Teharn, Iran³.