

Effect of systemic fungicides on total protein, carbohydrate and phenolic contents of *Solanum melongena* and *Avena sativa*

ZAMIN. S. SIDDIQUI

Department of Botany, University of Karachi, Karachi-75270, Pakistan

ABSTRACT

Effect of systemic fungicides viz. topsin-M (Methyl thiophenate) and benlate (benomyl) on protein, carbohydrate, and phenolic contents of *Solanum melongena* and *Avena sativa* were examined. Benlate (Benomyl) showed greater decrease in protein and carbohydrate contents of both the test species as compared to topsin-M (Methyl thiophenate). An increase in phenolic contents was recorded in the two species.

INTRODUCTION

The introduction of systemic fungicides is a major landmark in the history of chemical control of plant diseases. Topsin-M (Methyl thiophenate) and benlate (Benomyl) are systemic fungicides of benzimidazol group which are used for the control of diseases such as powdery mildew, downy mildew, brown rust, grey mould, rust and smuts of wheat (Thomson, 1985; Singh, 1991). Despite their use, systemic fungicides have been reported phytotoxic. The application of benomyl was found to produce chlorosis and irregular depression at the central and marginal portion of saffron leaves (Reyes, 1975). Alchor metaxyl induced sharp decrease in cell division (Coman et al., 1990). Bezimidazol-N-Sulfonamide interfered electron transport system (ETS) by combining NADH or succinate (Pillonel, 1993). Likewise carbendazim induced chromosomal aberration in somatic and germ cells of pearl and sunflower (Harchand et

al., 1991). It is presumed that the metabolic changes induced by fungicides will certainly affect the chemical composition of the host. This important aspect has been dealt only by a few workers (Berger and Cwick, 1990). The present study was undertaken to examine the effect of topsin-M (Methyl thiophenate) and benlate (benomyl) on *Avena sativa* and *Solanum melongena*. Protein and carbohydrate were selected as an indicator for changes in nutritive value and phenols as an indicator for the stress developed due to the presence of fungicides.

MATERIAL & METHODS

Seeds of *Avena sativa* and *Solanum melongena* obtained from National Institute of Agriculture and Biology (NIAB) Faisalabad were surface sterilized with 0.1% mercuric chloride for 10 min. Then washed with distilled deionized water.

The seeds were sown in 16" earthen pots containing 15 kg soil mixed with cowdung manure in 3:1 ratio. Ten seeds were sown in each pots. The pots were regularly watered and kept in glass house at temperature of 30-35°C and 35-45% R. H. Fifteen-day-old seedlings were sprayed separately with topsin-M (Methyl thiophenate) and benlate (benomyl). These fungicides were applied at the rate of 400, 600, 800 and 1000 mg/l with hand sprayer. the concentration of spray solution was based on formulations. Unsprayed plants were kept as control. Leaf samples were collected randomly after 2 weeks of treatment and changes in total protein content in leaves was measured after the extraction 5% cold trichloroacetic acid (TCA) and estimated through using folin phenol reagent, optical density read at 750 nm (Lowry *et al.*, 1951). Carbohydrate was detected by using anthrone reagent, optical density read at 520 nm (Yemm & Willis, 1956), while phenolic content was recorded after the extraction in ethanol and estimated using 10% folin reagent (Swain and Hillis, 1959), optical density was observed at 660nm. Erma photic 100 spectrophometer was used for each analysis and quantity expressed in mg/mg fresh weight by using standard curve. Three replicates were used for each analysis.

RESULT & DISCUSSION

Application of systemic fungicides viz. topsin-M (Methyl thiophenate) and benlate (benomyl) decreased the protein content of *S. melongena* and *A. sativa* (fig. 1). However, benlate (benomyl) exhibited phytotoxicity which resulted in greater reduction in protein content of *A. sativa* at 1000 mg/l, with reference to concentration of fungicide up and down trend was found. While slight increase in protein content was recorded in *Solanum melongena* when treated with benlate (benomyl) 800 mg/l.

Decrease in carbohydrate content of *S. melongena* and *A. sativa* was recorded (fig. 1). However, maximum decrease over control was measured in *A. sativa* treated with benlate (benomyl) 1000 mg/l.

An increase in phenolic content was observed in the two test species (fig. 1). However, maximum increase was recorded when plants were treated with benlate (benomyl) 1000 mg/l.

Application of systemic fungicide viz. benlate (benomyl) and topsin-M (Methyl thiophenate) have been increased during last 20-30 years. The metabolic changes induced as a result of phytotoxicity of systemic fungicides reduces mineral contents and inhibits biosynthesis of protein, carbohydrate and vitamins (Berger & Cwick, 1990). An osmotic shock effect of systemic fungicides results in the release of protein and loss of membrane transportability in the leaf cells (Amar & Rienhold, 1973). In contrast, in present study benlate (benomyl) has been found to increase NAD and NADP ratio (Godvary & Waywood, 1970) and increase in NADP and ATP level (Mishra & Waywood, 1968) by inducing a change in the enzyme system which may result in the conservation of leaf protein and chlorophyll in detached wheat leaves (Person *et al.*, 1957). It has been suggested that plants sprayed with systemic fungicides suffer from chemical stress (Ahmed & Siddiqi, 1995) and the compound produced as a result of stress may act as protective compound against pathogenic fungi (Friend, 1977; Ahmed & Siddiqi, 1995).

Stress condition causes abnormalities in the biochemical pathway due to which toxic phenolic compound like flavones are formed (Reld *et al.*, 1992). Compounds produced by chemical stress were potential inhibitors of germination and seedling growth (Heisy, 1990). Phytotoxin in the form of phenolic compounds are responsible for limiting cell




Fig. 1. Effect of topsin-M and benlate on protein, carbohydrate and phenolic contents of *Solanum melongena* and *Avena sativa*.

division, respiration, photosynthesis and disruption of cell membrane (Mucilas et al., 1992).

In the present studies benlate (benomyl) was found more toxic than topsin-M (Mehtyl thiophenate) a preferential use of topsin-M (Mehtyl thiophenate) over benlate (benomyl) may be suggested for plant disease control.

References

- AHMED, S. and Z. S. SIDDIQUI, 1995. Effect of Topsin-M (Methyl thiophenate) on chlorophyll, protein, and phenolic contents of *Hibiscus esculentus* and *Capsicum annum*. Pak. J. Bot. 27 (1): 175-178.
- AMAR, L. and L. REINHOLD. 1973. Loss of membrane transportability in leaf cell and release of protein as a result of osmotic shock. Plant Physiol., 51: 620-625.
- BERGER, S. and CWICK. 1990. Selected aspect of adverse nutritional effect of pesticides. Ernährung, 14 (7-8): 411-415.
- COMAN, N., M. DORDCA and t. PERSECA. 1990. The estimation of alcoholar and metaxylon cell division. Biologia, 35 (2): 55-60.
- FRIEND, J. 1977. Phenoilc substances and plant diseases. Recent adv. Phytochem., 12: 557.
- GODVARY, E. R. and WAYWOOD. 1970. Nicotinamide adenine dinucleo metabolism in plant, intermediate biosynthesis in wheat leaves and the effect of BZI. J. Bot. 48: 2267-2278.
- HARCHAND., R.; K. GROVER and P. K. SAREEN. 1991. Carbendazim induced chromosomal aberration in somatic and reproduction of pearl and sunflower. Cycologia, 56(4): 559-566.
- HEISY, R. M. 1991. Allelopathic and herbicidal effect of extract from tree of heaven (*Althaus altissima*) Amar. J. Bot, 77 (5): 230-235.
- LOWRY, O. H.; N. J. ROSEBROUGH; A. L. PARR and N. J. RANDALL, 1951. Protein measurment with Folin Phenol reagent. J. Biol. Chem., 193: 265-275.
- MACIAS, S. A., J. C. G. GALINDO and G. M. MASSANOT. 1992. Potential allelopathic of several sesquiterpene Lactone models. Phytochem., 31: 1969-1977.
- MISHRA, D. and E. R. WAYWOOD. 1968. The effect of BZI and Kinetin on the nicotinamide and nucleotide content of senescence wheat leaf. Can. J. Bio., 46:

167-178.

- PRESON, C. D.; J. SAMBROSKI and F. R. FORSYTH. 1957. The effect of BZI on detached wheat leaves. *Can. J. Bot.*, 180: 1294-1295.
- PILLONEL, C. 1993. Interaction of benzimidazol-N-sulfonamide with the cytochrome b/c complex *Phythium alphanidermatum*. *Pesticide Sci.*, 43(2): 107-113.
- REID, L. M.; D. E. MATHER.; J. AMASON.; T. HAMILTON and R. J. BOLTON. 1992. Changes in phenolic constituent of maize silk infected with *Fusarium graminearum*. *Can. j. Bot.*, 70: 1697-1700.
- REYES, A. A. 1975. Phytotoxicity of benomyl to saffron. *Phytopathology.*, 65: 1-6.
- SINGH, R. S. 1991. Management of disease with chemical. In: Introduction to principle of plant pathology. 3rd edition. Oxford & IBH Pub. New Delhi. India.
- SWAIN, T. and W. E. HILLIS. 1959. The phenolic constituent of *Prunus domestica*. *J. Sci. Food. Agric.* 10: 63-68.
- THOMSON, W. T. 1985. Agricultural chemical. Book 4 Fungicide. Thomson Pub. USA. pp. 126-135.
- YEMM, F. W. and A. J. WILLIS. 1956. The estimation of carbohydrate in plant extracts by anthrone. *J. Biochem.*, 57: 508.

Address of the author: Dr. ZAMIN. S. SIDDIQUI, Department of Botany, Unversity of Karachi, Karachi-75270, Pakistan.