

Ent. Phyt. Appliq .
Vol . 53 , No . 1 &2, F  v . 1986

POSSIBILITY OF CODLING MOTH CONTROL BY
INSECTICIDAL MEASURES IN ESFAHAN¹

N. DASTGHEYB BEHESHTI²

SUMMARY

Four and five sprays were applied against the three generations of codling moth (*Laspeyresia pomonella* L). to indicate the degree of effectiveness of insecticidal measures.

In both treatments, the first generation was treated twice and the other sprays were used against 2nd and 3rd generations (See Figure 1).

There has been no significant difference between four and five times of insecticide application while both have been found to be quite effective against codling moth.

REFERENCES

- RADJABI, GH., BEHESHTI, N. D., AKRAMI, F. & BAYATASSADI, H., 1977. Codling moth in Iran, researches made from 1975 - 1977. Ent. Phyt. Appl., Vol. 46, No.1 &2 (in Farsi with summary in English).
- RADJABI, GH. BEHESHTI, N. D., AKRAMI, F. and BAYATASSADI, H., 1979. Possibility of population reduction of Codling moth with the control of the first generation in Iran. Ent. Phyt. Appl. Vol. 48, No. 1 (in Farsi with a summary in English).

1 - Received for publication, Jan. 1, 1985.

2- Eng. Nahid Dastgheyb Beheshti, Plant Pests and Diseases Research Laboratory, P. O. Box 419, Esfahan, Iran.